

FLOWERING SEASONS OF AUSTRALIAN PLANTS.

By E. HAVILAND, F.L.S.

No. 3.—PLANTS FLOWERING IN THE NEIGHBOURHOOD OF SYDNEY DURING THE MONTH OF SEPTEMBER, IN ADDITION TO THOSE ENUMERATED IN LISTS FOR JULY AND AUGUST, MOST OF WHICH ARE STILL IN FLOWER.

Dilleniaceæ—	Ficoideæ—
<i>Hibbertia fasciculata.</i>	<i>Tetragonia expansa.</i>
Cruciferae—	Leguminosæ—
<i>Cakile maritima.</i>	<i>Bossicea microphylla</i>
Violaceæ—	<i>Acacia stricta</i>
<i>Viola betonicifolia.</i>	<i>Pultencea flexilis</i>
Pittosporæ—	„ <i>retusa</i>
<i>Marianthus procumbens</i>	<i>Daviesia ulicina</i>
<i>Pittosporum undulatum.</i>	<i>Gompholobium minus</i>
Polygalæ—	„ <i>latifolium</i>
<i>Comesperma volubile.</i>	<i>Mirbelia reticulata.</i>
Tremandreae—	Myrtaceæ—
<i>Tetratheca juncea</i>	<i>Leptospermum parvifolium</i>
„ <i>ericoides.</i>	„ <i>attenuatum</i>
Rutaceæ—	<i>Bæckea crenulata</i>
<i>Correa speciosa</i> var <i>canensis</i>	„ <i>diosmifolia.</i>
<i>Philotheca australis.</i>	Rhamnaceæ—
Linææ—	<i>Pomaderris lanigera</i>
<i>Linum marginale.</i>	„ <i>elliptica</i>
Euphorbiaceæ—	Loranthaceæ—
<i>Phyllanthus thymoides</i>	<i>Notothixos subaureus.</i>
<i>Breynia oblongifolia.</i>	Proteaceæ—
Stackhousiaceæ—	<i>Grevillea laurifolia</i>
<i>Stackhousia muricata</i>	„ <i>buxifolia</i>
„ <i>viminea.</i>	<i>Hakea dactyloides.</i>

Compositæ—

Helichrysum diosmifolium.

Goodeniaceæ—

Goodenia barbata.

Plantagineæ—

Plantago debilis.

Solanaceæ—

Solanum aviculare.

Scrophularinæ—

Veronica plebeia.

Labiataæ—

Plectranthus parviflorus.

Epacrideæ—

*Styphelia triflora**Monotoca scoparia.*

Orchideæ—

Caladenia testacea,, *alba**Thelymitra ixioides**Diuris elongata*,, *punctata**Caleana major**Calochilus paludosus**Glossodia minor**Prasophyllum elatum.*

Irideæ—

Patersonia glabrata.

Amaryllideæ—

Hypoxis hygrometrica.

Liliaceæ—

Burchardia umbellata.